arXiv:1207.3602v2 [hep-th] 24 Aug 2012

Prepared for submission to JHEP

Twistor-strings and gravity tree amplitudes

Tim Adamo and Lionel Mason

The Mathematical Institute
University of Oxford
24-29 St. Giles', Oxford OX1 3LB
United Kingdom

E-mail: `adamo@maths.ox.ac.uk`, `lmason@maths.ox.ac.uk`

Abstract: Recently we proposed that the original Berkovits-Witten twistor-string theory could be used to compute Einstein supergravity tree amplitudes. Subsequently, remarkable formulae for the MHV amplitudes were found by Hodges and generalized to N^{k}MHV by Cachazo and Skinner. Here we show how the Hodges matrix and its higher MHV-degree generalization can be understood as arising from the twistor-string formula. In particular we see that if all contractions in the worldsheet correlator calculations are allowed, the wrong answers are obtained. On the other hand we obtain the correct answer if the contractions form connected trees (alternatively, the contractions in the correlators must leave a minimal number of disconnected terms). The generalised Hodges matrix arise as weighted Laplacian matrices for the graph of possible contractions. The reduced determinants of these weighted Laplacian matrices give the sum of of the connected tree contributions by the Matrix-Tree theorem.

Contents

1 Introduction **1**

2 The Twistor-string Formula for Conformal and Einstein Gravity **4**
- 2.1 Twistor-strings for conformal supergravity 4
- 2.2 Reduction to Einstein gravity 6

3 Evaluating the Twistor-string Tree Formula **8**
- 3.1 Dependence on Λ 8
- 3.2 Naïve computation of the $O(\Lambda^0)$ contribution 8
- 3.3 Connected trees and Feynman diagrams 9
- 3.4 The $O(\Lambda^0)$ case 10

4 The MHV Amplitude From the $O(\Lambda)$ Contribution **13**
- 4.1 The 3-point amplitude 16

5 The N^{k}MHV Amplitude **17**

6 Discussion **19**

A Graph Theory **20**

1 Introduction

The twistor-string introduced by Witten [1] and realized as a momentum space tree formula [2] for $\mathcal{N} = 4$ super-Yang-Mills has been a remarkable stimulus for developments in our understanding of gauge theory amplitudes (see [3] for a review biased towards the interests of these authors and [4] and other reviews in that volume for other influences). The question naturally arises as to whether analogous ideas can be made to work for gravity. Witten's twistor-string theory does contain conformal gravity, a fourth order conformally invariant theory whose action is given by the square of the Weyl tensor. However, at this point no twistor-string theory is known that produces just Einstein gravity, although the recent formulae of Cachazo and Skinner [5, 6] (see also [7, 8]) for $\mathcal{N} = 8$ supergravity amplitudes from rational curves in twistor space are remarkably suggestive of the existence of such a theory.

In [9] it was argued, using an observation of Maldacena [10], that the twistor-string tree formula for conformal supergravity amplitudes due to Berkovits and Witten [11] could be used to calculate Einstein tree amplitudes. The procedure yields an extra factor of the cosmological constant Λ and hence should vanish as $\Lambda \to 0$. However, for an n-particle amplitude, the procedure automatically yields a polynomial of degree-n in Λ so that it is straightforward to divide by Λ and take the limit $\Lambda \to 0$. In [9], only three-point MHV and anti-MHV amplitudes were checked in momentum space (found for all Λ and checked in the $\Lambda \to 0$ limit). In this article we investigate more general formulae that are yielded in momentum space in the $\Lambda = 0$ limit.

The first thing we observe is that the formulae in [9] must be understood in a semi-classical context in which only connected trees[1] are allowed amongst the contractions in the correlator in order to obtain the correct answer when $\Lambda = 0$. Indeed, the answer is simply non-vanishing at $\Lambda = 0$ if loops and more disconnected terms amongst the contractions in the correlation functions are allowed.

We move on to consider the $O(\Lambda)$ part of the conformal gravity amplitude, which by the Maldacena argument above yields the Einstein gravity contribution at $\Lambda = 0$. Working with just connected trees, we see that the tree formulae for the MHV amplitude of [12] can be identified simply with the Feynman diagrams of contractions required for the worldsheet correlators restricted to being connected trees. Furthermore, by summing the diagrams using a weighted extension of the Matrix-Tree theorem, we can obtain a more fundamental understanding of the origin of Hodges' recent remarkable MHV formula [13], as well as one of the reduced determinant factors in the N^{k}MHV formula of Cachazo and Skinner [5, 6]. These Hodges matrices can be understood as weighted Laplacian matrices for all possible contractions extended to a permutation invariant framework. The reduced determinants that arise in these formulae come from extensions of the Matrix-Tree theorem. We therefore obtain not only a more fundamental explanation for the origin of the ingredients in Hodges' formulae, but also support for the assertions of [9] for MHV, at least at $O(\Lambda)$ and with the proviso on the restrictions of allowable contractions in the twistor-string correlation functions.

The conformal gravity twistor-string formulae are very much rooted in $\mathcal{N} = 4$ supersymmetry and so do not manifest full permutation symmetry. However, if they are correct, they must have an emergent permutation symmetry presumably arising from a $\mathcal{N} = 8$ formulation of the momentum space analogue of the tree-formula. As already mentioned, such a formula has independently been found by Cachazo and Skinner [5, 6]. In this paper we see how the Hodges formula and certain key ingredients of the Cachazo-Skinner formula naturally arise from the Berkovits-Witten twistor-string theory. We obtain a fairly complete picture at MHV but rather less complete at higher MHV-degree.

We now give a brief review of the Cachazo-Skinner formula. It is a natural extension of Hodges' formula for the MHV amplitude, and is a function of the kinematic invariants of the momenta $\lambda_i\tilde{\lambda}_i$ where $i = 1, \ldots, n$ index the external particles and λ_i is a two-component Weyl spinor λ_{iA}, $A = 0, 1$. For $\tilde{\lambda}_i$, we understand it to also include the fermionic momenta so that $\tilde{\lambda}_i = (\tilde{\lambda}_{iA'}, \eta_i^a)$ where $A' = 0'1'$ are spinor indices and $a = 1, \ldots 8$ is a R-symmetry index. It also uses auxilliary variables $\sigma_i = (\sigma_{i0}, \sigma_{i1})$ that are homogeneous coordinates for n marked points on the Riemann sphere (the string worldsheet). We use square brackets for primed spinor contractions, angle brackets for unprimed spinor contractions and round brackets for contractions of the σs.

---

[1] Rather than ruling out loops, we can require simply that the number of disconnected components amongst the contractions is minimal; we thank Mat Bullimore for this observation.

Following [5, 6, 13], we introduce a $n \times n$ matrix function of the kinematic invariants and σ_is at N^{k}MHV degree (R-charge $k+2$) by

$$\tilde{\phi}^k_{ij} = \begin{cases} \frac{[ij]}{(i\,j)}\left(\frac{(\xi\; j)}{(\xi\; i)}\right)^{k+2} & i \neq j \\ -\sum_{l\neq i} \frac{[i\,l]}{(i\,l)}\left(\frac{(\xi l)}{(\xi i)}\right)^{k+2} & i = j \end{cases}, \tag{1.1}$$

where ξ is an arbitrary fixed point on $\mathbb{CP}^1$. The off-diagonal entries of $\tilde{\phi}^k$ are just the twistor-string propagators associated to contractions among worldsheet vertex operators. This is related to the matrix $\tilde{\Phi}^k$ appearing in [5, 6] (and in [13] for $k = 0$) via conjugation by the matrix $T = \operatorname{diag}\left((\xi\; i)^{k+2}\right)$:

$$\tilde{\Phi}^k_{ij} = \begin{cases} \frac{[ij]}{(i\,j)} & i \neq j \\ -\sum_{l\neq i} \frac{[i\,l]}{(i\,l)}\left(\frac{(\xi l)}{(\xi i)}\right)^{k+2} & i = j \end{cases}, \qquad \tilde{\phi}^k = T^{-1}\tilde{\Phi}^k T. \tag{1.2}$$

Hence, reduced determinants of $\tilde{\phi}^k$ and $\tilde{\Phi}^k$ will be equivalent[2].

When $k = 0$, (1.2) is Hodges' matrix for the MHV formula, which has three-dimensional kernel as a consequence of momentum conservation. The general $\tilde{\Phi}^k$ has a $k+3$-dimensional kernel given by the relations

$$\sum_{j=1}^{n} \tilde{\Phi}^k_{ij}\sigma_j^{A_1}\cdots\sigma_j^{A_{k+2}} = 0\,, \tag{1.3}$$

which follow from the support of delta functions in the formula. A type of Fadeev-Popov argument leads to an invariant determinant $\det'(\tilde{\Phi}^k)$ of this matrix generalizing that described by Hodges. This is the main new ingredient in the tree formula beyond the Yang-Mills case. The final formula can be expressed as

$$\mathcal{M}^k_n(1,\ldots,n) = \int \frac{\prod_{r=0}^{k+1}\mathrm{d}^{4|8}U_r}{\mathrm{vol}\;\mathrm{GL}(2,\mathbb{C})}\det{}'(\tilde{\Phi}^k)\det{}'(\Phi^k)\prod_{i=1}^{n}\mathrm{D}\sigma_i\; h_i\left(\lambda_i\tilde{\lambda}_i; Z(\sigma_i)\right), \tag{1.4}$$

where the U_r are parameters for the degree $k+1$ map $Z : \mathbb{CP}^1 \to \mathbb{PT}$; $\det'(\Phi^k)$ and $\det'(\tilde{\Phi}^k)$ are modified determinants; and the h_i are momentum eigenstates.

It is instructive to compare this to the analogous formula for $\mathcal{N} = 4$ Yang-Mills:

$$\mathcal{A}^k_n(1,\ldots,n) = \int \frac{\prod_{r=0}^{k+1}\mathrm{d}^{4|4}U_r}{\mathrm{vol}\;\mathrm{GL}(2,\mathbb{C})}\prod_{i=1}^{n}\frac{\mathrm{D}\sigma_i}{(i\; i+1)}a_i\left(\lambda_i\tilde{\lambda}_i; Z(\sigma_i)\right). \tag{1.5}$$

As remarked in [2], after integrating out part of the moduli space, both formulae have as many delta functions as integrals and the result reduces to a sum of residues at the solutions (both real and complex) of the equations implied by delta-functions. The main difference between (1.4) and (1.5) is that the Parke-Taylor-like denominator of $\mathcal{N} = 4$ is replaced by the product of determinants $\det'(\tilde{\Phi}^k)\det'(\Phi^k)$.

---

[2] The fact that (1.2) and hence (1.1) is independent of the choice $\xi \in \mathbb{CP}^1$ follows from the delta-function support of the amplitude; see [5, 6] for more details.

This paper gives an interpretation of the $\det'(\tilde{\Phi}^k)$ factor in (1.4) from twistor-string theory. Indeed, the individual off-diagonal entries of the matrix are precisely all the possible propagators arising from a certain type of contraction in the world-sheet conformal field theory extended to a permutation invariant framework. The diagonal entries are then precisely what is required to obtain the weighted Laplacian matrix for the graph of all possible contractions. We will see that at N^{k}MHV, only $n-3-k$ such contractions should be allowed and the reduced determinant is constructed precisely as the $n-3-k$ minor that effectively sums trees with just $n-3-k$ such propagators.

There are nevertheless significant gaps in our understanding here. In particular, we do not yet understand how the twistor-string ingredients can be used to assemble $\det'\Phi$, nor the Vandermonde factors in the definition of $\det'\tilde{\Phi}$. If the Berkovits-Witten twistor-string theory is to correctly describes tree-level conformal gravity, it must also be possible to obtain these within that framework. Indeed it is so closely connected to the twistor-action for conformal gravity [14], which has been systematically shown to give rise directly to classical conformal gravity, that it seems unlikely it will not be possible to obtain these ingredients, but a deeper understanding is required. Of course, of independent interest, the considerations in this paper are also likely to play a role in any yet to be understood $\mathcal{N} = 8$ twistor-string theory for Einstein gravity.

2 The Twistor-string Formula for Conformal and Einstein Gravity

Here we briefly summarize the ingredients used in our twistor-string construction; more details can be found in [9]. Note that we abuse notation slightly, since in this section there is a $\mathrm{SU}(4)_R$ R-symmetry associated with the twistor-string construction. We will use similar notation for both this $\mathcal{N} = 4$ and the $\mathcal{N} = 8$ R-symmetry used elsewhere.

2.1 Twistor-strings for conformal supergravity

Non-projective twistor space is $\mathbb{T} = \mathbb{C}^{4|4}$ and projective twistor space is $\mathbb{PT} \cong \mathbb{CP}^{3|4} = \mathbb{T}/\{Z \sim \mathrm{e}^{\alpha} Z\}$, for $\alpha \in \mathbb{C}$. A twistor will be represented as $Z^I \in \mathbb{T}$, $Z^I = (Z^\alpha, \chi^a)$, $\alpha = 0, \dots 3$, $a = 1, \dots, 4$ with Z^α bosonic and χ^a fermionic and the bosonic part $Z^\alpha = (\lambda_A, \mu^{A'})$, $A = 0, 1$, $A' = 0'1'$. A point $(x, \theta) = (x^{AA'}, \theta^{Aa})$ in chiral super Minkowski space-time $\mathbb{M}$ corresponds to the $\mathbb{CP}^1$ (complex line) $X \subset \mathbb{PT}$ via the incidence relation

$$\mu^{A'} = i x^{AA'} \lambda_A \,, \quad \chi^a = \theta^{Aa} \lambda_A \,. \tag{2.1}$$

with λ_A homogeneous coordinates along X.

We use a closed string version of the Berkovits model[3] [11, 16] with a Euclidean world-sheet Σ. The fields are

$$Z : \Sigma \to \mathbb{T} \,, \quad Y \in \Omega^{1,0}(\Sigma) \otimes T^*\mathbb{T} \,, \quad \text{and} \quad a \in \Omega^{0,1}(\Sigma) \,.$$

---

[3] The standard Berkovits open twistor-string would be just as good for most of the considerations in this paper; although it is tied into split signature and gets some signs wrong, subtleties over choices of contours are avoided. For more on the version used here see [15].

The action is

$$S[Z,Y,a] = \int_\Sigma Y_I \bar\partial Z^I + a Z^I\, Y_I\,, \tag{2.2}$$

up to matter terms, and has the gauge freedom

$$(Z,Y,a) \to (\mathrm{e}^{\alpha} Z, \mathrm{e}^{-\alpha} Y, a - \bar\partial\alpha), \qquad \alpha \in \mathbb{C}.$$

The gauging reduces the string theory to one in $\mathbb{PT}$ and the formalism allows one to use homogeneous coordinates on $\mathbb{PT}$.

Amplitudes are computed as an integral of worldsheet correlators of vertex operators on Σ over the moduli space of 'zero-modes': the space of classical solutions to the equations of motion. For gravity, the vertex operators correspond to deformations of the complex structure together with the cohomology class of deformations of the B-field. These are given by $\bar\partial$-closed $(0,1)$-forms F on the bosonic part of twistor space $\mathbb{PT}$ with values in the generalized tangent space $T\oplus T^*\mathbb{PT}$. On $\mathbb{T}$ we represent these by $\bar\partial$-closed $(0,1)$-forms $F := (f^I, g_I)$ of homogeneity $(1,-1)$ satisfying $\partial_I f^I = 0 = Z^I g_I$, defined modulo gauge transformations $(\alpha Z^I, \partial_I \beta)$. These conditions imply that $(f^I\partial_I, g_I \mathrm{d}Z^I)$ represents a section of $T\oplus T^*\mathbb{PT}$. The corresponding vertex operators take the form

$$V_F := V_f + V_g := \int_\Sigma f(Z)^I Y_I + g(Z)_I \mathrm{d}Z^I\,.$$

For n-particle tree-level amplitudes, we take $\Sigma \cong \mathbb{CP}^1$ and the amplitude reduces to an integral of a correlation function of n vertex operators

$$\mathcal{M}(1,\ldots,n) = \sum_{d=0}^{\infty} \int_{\mathscr{M}_{d,n}} \mathrm{d}\mu_d \langle V_{F_1} \ldots V_{F_n}\rangle_d\,, \tag{2.3}$$

over the space $\mathscr{M}_{d,n}$ of maps $Z : \mathbb{CP}^1 \to \mathbb{PT}$ of degree-d and n marked points.[4] This is the moduli space of zero-modes for the twistor-string, and has a mathematical definition as a supersymmetric generalization of Kontsevich's moduli space of stable maps [17]. To be more concrete, we can represent the maps by

$$Z(\sigma) = \sum_{r=0}^{d} \sigma_0^r \sigma_1^{d-r} U_r\,, \quad \mathrm{d}\mu_d = \frac{1}{\mathrm{vol}\mathrm{GL}(2,\mathbb{C})} \prod_{r=0}^{d} \mathrm{d}^{4|4} U_r\,, \tag{2.4}$$

where σ_A are homogeneous coordinates on $\mathbb{CP}^1$ and $U_r \in \mathbb{T}$ provide a set of coordinates on $\mathscr{M}_{d,0}$ with redundancy $\mathrm{GL}(2,\mathbb{C})$ acting on the σ and hence the U_r. The vertex operator $V_{F_i} = V_F(Z(\sigma_i))$ is inserted at the ith marked point $\sigma_i \in \Sigma$, and the correlator naturally introduces a $(1,0)$-form at each marked point either from the Y_I or the $\mathrm{d}Z^I$, whereas the 'wave-functions' (f^I, g_I) naturally restrict to give a $(0,1)$-form at each marked point. See [18] for further explanation.

---

[4] The rules for taking the correlators are different at different degrees, hence the subscript d on the correlator.

The correlators are computed by performing Wick contractions of all the Ys with Zs to give the propagator

$$\langle Y(\sigma)_I Z^J(\sigma')\rangle_d = \left(\frac{(\xi\sigma')}{(\xi\sigma)}\right)^{d+1} \frac{\delta_I^J \mathrm{D}\sigma}{(\sigma\sigma')}\,, \quad \mathrm{D}\sigma = (\sigma\; \mathrm{d}\sigma)\,, \quad (\sigma\;\sigma') = \sigma_0\sigma_1' - \sigma_1\sigma_0'\,, \qquad (2.5)$$

where $(\sigma\sigma') = \epsilon_{AB}\sigma^A\sigma'^B$ is the $\mathrm{SL}(2,\mathbb{C})$ invariant inner product on the world-sheet coordinates. When Y acts on a function of Z at degree d, it then differentiates before applying the contraction; Y acting on the vacuum gives zero so that all available Ys must be contracted, but this contraction can occur with any available Z. The ξ is an arbitrary point on the Riemann sphere and reflects the ambiguity in inverting the $\bar\partial$-operator on functions of weight d on Σ. The overall formula should end up being independent of the choice of ξ.

Unlike the Yang-Mills case, the degree d of the map is not directly related to the MHV degree of the amplitude (which essentially counts the number of negative helicity gravitons minus 2). The MHV degree of an amplitude counts the number of insertions of V_g minus 2 (so the MHV amplitude has two V_gs). Conformal supergravity amplitudes have been calculated from this formula in [19, 20].

2.2 Reduction to Einstein gravity

We use a modified form of the Maldacena argument [10] in which the cosmological constant is not normalized, so that we can investigate the limit $\Lambda \to 0$. The argument then shows that the classical S-matrix for conformal supergravity (i.e., the action evaluated perturbatively on in- and out-wave functions) agrees with that for Einstein gravity, but multiplied by an additional factor of the cosmological constant Λ.

Even for Einstein gravity without supersymmetry, we will need to use the supergeometry of $\mathcal{N}=4$ supertwistor space.[5] We first give the restriction required of the vertex operators for $\mathcal{N}=4$ supersymmetry and then for $\mathcal{N}=0$.

To reduce to the Einstein case we must break conformal invariance. This is done by introducing skew infinity twistors I_{IJ}, I^{IJ} with super-indices $I, J = (\alpha, a)$. The bosonic parts $I_{\alpha\beta}$, $I^{\alpha\beta}$ satisfy

$$I^{\alpha\beta} = \frac{1}{2}\varepsilon^{\alpha\beta\gamma\delta} I_{\gamma\delta}\,, \quad I^{\alpha\beta} I_{\beta\gamma} = \Lambda\delta^\alpha_\gamma\,,$$

where Λ is the cosmological constant and we shall set the fermionic part equal to zero. In terms of the spinor decomposition of a twistor $Z^\alpha = (\lambda_A, \mu^{A'})$ we have

$$I_{\alpha\beta} = \begin{pmatrix} \varepsilon^{AB} & 0 \\ 0 & \Lambda\varepsilon_{A'B'} \end{pmatrix}\,, \qquad I^{\alpha\beta} = \begin{pmatrix} \Lambda\varepsilon_{AB} & 0 \\ 0 & \varepsilon^{A'B'} \end{pmatrix}\,. \qquad (2.6)$$

They have rank two when $\Lambda = 0$ (i.e., the cosmological constant vanishes) and four otherwise. The fermionic parts of I^{IJ} can be non-zero and correspond to a gauging for the R-symmetry of the supergravity [21, 22].

---

[5] At tree-level we can pull out the pure gravity parts of the amplitude, but their construction still relies on the fermionic integration built into the twistor-string formulae; this simply leads to some spinor contractions.

Geometrically I^{IJ} and I_{IJ} respectively define a Poisson structure $\{,\}$ of weight -2 and contact structure τ of weight 2 by

$$\{h_1, h_2\} := I^{IJ}\partial_I h_1 \partial_J h_2 \,, \qquad \tau = I_{IJ} Z^I \mathrm{d}Z^J \,,$$

and we can use the Poisson structure to define Hamiltonian vector fields $X_h = I^{IJ}\partial_I h \partial_J$ which are homogeneous when h has weight 2. Lines in $\mathbb{PT}$ on which the contact form τ vanishes correspond to points at infinity. This zero-set defines a surface $\mathscr{I}$ in space-time which is null when $\Lambda = 0$, space-like for $\Lambda > 0$, and time-like for $\Lambda < 0$.

The Einstein vertex operators $(V_h, V_{\tilde{h}})$ correspond to $V_f + V_g$ subject to the restriction

$$(f^I, g_I) = (I^{IJ}\partial_J h, \tilde{h} I_{IJ} Z^J)$$

so that

$$V_h = \int_\Sigma I^{IJ} Y_I \partial_J h \,, \quad V_{\tilde{h}} = \int_\Sigma \tilde{h} \wedge \tau \,. \tag{2.7}$$

The first of these gives the $\mathcal{N} = 4$ multiplet containing the negative helicity graviton and the second that containing the positive helicity graviton. See [9] for more discussion. In order to reduce to standard non-supersymmetric Einstein gravity we must impose

$$h = e_2 \,, \qquad \text{and} \qquad \tilde{h} = \chi^4 \tilde{e}_{-6} \,. \tag{2.8}$$

Thus, evaluated on vertex operators constructed from (2.7) with (2.8), (2.3) leads to the construction of Einstein gravity tree amplitudes. With this restriction, we see that there is now a correlation between degree of the maps and MHV degree, as fermionic variables only come with $\tilde{h}$: since there are $4d$ fermionic integrations in the path integral for the amplitude, there must be d insertions of $\tilde{h}$.

At $\mathcal{N} = 4$ there are spurious amplitudes that can be constructed from the Einstein wave functions corresponding to other conformal supergravity sectors. In [9], we argued that the Einstein gravity amplitudes could be isolated by imposing the correspondence between the degree of the map and the MHV degree: $d = k + 1$. This leads to the following starting point for tree-level scattering amplitudes in Einstein gravity from twistor-string theory:

$$\begin{aligned}
\mathcal{M}_n^k(1, \ldots, n) &= \lim_{\Lambda \to 0} \frac{1}{\Lambda} \int_{\mathscr{M}_{k+1,n}} \mathrm{d}\mu_{k+1} \left\langle V_{h_1} \cdots V_{h_{n-k-3}} V_{\tilde{h}_{n-k-2}} \cdots V_{\tilde{h}_n} \right\rangle_{k+1} \\
&= \lim_{\Lambda \to 0} \frac{1}{\Lambda} \int_{\mathscr{M}_{k+1,n}} \mathrm{d}\mu_{k+1} \left\langle Y \cdot \partial h_1 \cdots Y \cdot \partial h_{n-k-3} \; \tilde{h}_{n-k-2} \tau_{n-k-2} \cdots \tilde{h}_n \tau_n \right\rangle_{k+1} \\
&= \lim_{\Lambda \to 0} \frac{1}{\Lambda} \int_{\mathscr{M}_{k+1,n}} \mathrm{d}\mu_{k+1} \, \mathcal{C}_{n,k+1} \,,
\end{aligned} \tag{2.9}$$

where $Y \cdot \partial h_i = I^{IJ} Y_{I\, i} \partial_{J\, i} h_i$ and the last equation defines $\mathcal{C}_{n,k+1}$ as the worldsheet correlation function of the relevant vertex operators.

In this paper we will focus on the $\Lambda \to 0$ limit, in particular on the $O(\Lambda^0)$ and $O(\Lambda)$ parts. However, according to the Maldacena argument, if the Berkovits-Witten formulation correctly gives conformal supergravity amplitudes, then (2.9) will do so for Einstein gravity for all Λ.

3 Evaluating the Twistor-string Tree Formula

We now turn to the evaluation of the formula (2.9) to test the claim that it will produce Einstein gravity amplitudes correctly. The most elementary consequence is that at $\Lambda = 0$, $\mathcal{C}_{n,k+1}$ must vanish because of the overall extra factor of Λ. We will also be interested in the coefficient of Λ since according to the version of the Maldacena argument, this will give the $\Lambda = 0$ Einstein amplitudes which are now well known.

3.1 Dependence on Λ

Powers of Λ arise from the identity $I^{\alpha\beta} I_{\beta\gamma} = \Lambda \delta^{\alpha}_{\gamma}$. This gives a factor of Λ whenever there is a Wick contraction between a Y which is always be attached to an upstairs infinity twistor and a τ which contains a downstairs infinity twistor as follows:

$$\begin{aligned}\left\langle I^{\alpha\beta} Y_\beta(\sigma)\, \tau(\sigma')\right\rangle &= I^{\alpha\beta} \left\langle Y_\beta(\sigma)\, Z^\gamma(\sigma')\, \partial' Z^\delta(\sigma')\right\rangle\, I_{\gamma\delta} \\ &= I^{\alpha\beta} I_{\gamma\delta} \frac{\mathrm{D}\sigma}{(\xi\, \sigma)^{k+2}} \left(\delta^\gamma_\beta \frac{(\xi\sigma')^{k+2}}{(\sigma\sigma')} \partial' Z^\delta + \delta^\delta_\beta Z^\gamma \partial' \frac{(\xi\, \sigma')^{k+2}}{(\sigma\sigma')} \right) \\ &= \Lambda \frac{\mathrm{D}\sigma}{(\xi\, \sigma)^{k+2}} \left(\frac{(\xi\sigma')^{k+2}}{(\sigma\sigma')} \partial' Z^\alpha - Z^\alpha \partial' \frac{(\xi\, \sigma')^{k+2}}{(\sigma\sigma')} \right) ,\end{aligned}$$

where ∂' is the holomorphic exterior derivative with respect to σ'. Thus, at $O(\Lambda^0)$ we must exclude all Wick contractions with a τ. At $O(\Lambda)$ we are allowed one such contraction and indeed we will see that the non-trivial contribution has precisely one such τ-contraction.

3.2 Naïve computation of the $O(\Lambda^0)$ contribution

At $O(\Lambda^0)$ we exclude any contraction of a Y_i with a τ, so a Y_i can only contract with one of the wave functions. In the naïve calculation we will allow all such contractions. However, we will see that this does not vanish even for the MHV amplitude.

Working at $k = 0$ in (2.9), insert the standard momentum eigenstates

$$h_j = \int_{\mathbb{C}} \frac{\mathrm{d}s_j}{s_j^3} \bar{\delta}^2(s_j \lambda_j - p_j) \exp\left(i s_j [[\mu_j \tilde{\lambda}_j]] \right) . \tag{3.1}$$

In the $\Lambda \to 0$ limit we find

$$\langle I^{IJ} Y_{I\, i} \partial_J h_i\; h_j \rangle = \frac{[ij]}{\langle ij \rangle} \frac{\langle \xi j \rangle^2}{\langle \xi i \rangle^2} h_i h_j .$$

After performing all world-sheet integrals, the naïve contribution from the ith Y_i to the $\Lambda = 0$ part of the correlator is:

$$\sum_{j \neq i} \frac{[ij]}{\langle ij \rangle} \frac{\langle \xi j \rangle^2}{\langle \xi i \rangle^2} \equiv -\phi^i_i ,$$

borrowing Hodges' notation [13]. Hodges shows that it follows from momentum conservation that this is independent of ξ and indeed has an interpretation as an inverse soft factor for inserting a particle at the site i. This gives

$$\mathcal{C}_{n,1}|_{O(\Lambda^0)} = \prod_{i=1}^{n-2} -\phi^i_i ,$$

but this does not generically vanish.

3.3 Connected trees and Feynman diagrams

Rather than give up and deduce that the Berkovits-Witten twistor-string is incorrect, we instead use this as an indication that a different rule is needed for evaluating the twistor-string formula. The rule we will impose is that the Feynman diagrams for the contractions in the correlators appearing in (2.9) must be connected trees. This eliminates the loops and minimises the number of disconnected terms which are implicitly included in the naïve calculation above. We will see that it leads to the Hodges [13] formula at MHV via the weighted Matrix-Tree theorem, an analogue of Kirchoff's theorem for directed weighted graphs (see appendix A and [23–25]). It also sheds light on the possible twistor-string origin of Cachazo and Skinner's recent formula for tree-level N^{k}MHV amplitudes [5, 6]. As a bonus, it also provides an understanding of the origins for the tree formulae of [12] and [26] in twistor-string theory.

Our argument represents terms contributing to the correlators of (2.9) as Feynman diagrams of the world-sheet CFT. It will be useful to represent such diagrams graphically, so we set out our notation here. There are two different kinds of vertices and two different kinds of propagator (or contraction) which can contribute to $\mathcal{C}_{n,k+1}$. Vertices correspond to either $Y_i \cdot \partial h_i$ (white) or $\tilde{h}_j \tau_j$ (black). Straight solid directed edges correspond to propagators arising from contractions directed outwards from a Y_i to an Einstein wave function (either h_j or $\tilde{h}_j$); this utilizes the Poisson structure of the twistor space. Straight dashed edges correspond to a contraction directed outwards from a Y_i to a τ_j, and involve the contact structure. These both arise from standard contractions in the Berkovits-Witten twistor-string theory [11, 18]. See figure 1. We will only allow contributions from diagrams that are trees (i.e., no closed cycles) and connected *after removing all black vertices.*


i
$\tilde{h}_i \tau_i$
i
j
$\langle Y_i h_j \rangle = \frac{[ij]}{(ij)} \left(\frac{(\xi\, j)}{(\xi\, i)} \right)^{k+2}$
i
j
$\langle Y_i \tau_j \rangle$
i
$Y_i \cdot \partial h_i$

Figure 1. *Building blocks for Feynman diagrams*

The diagrams acquire a factor of Λ^r if there are r $\langle Y_i \tau_j \rangle$ contractions and we will just consider the cases where $r = 0$ or $r = 1$.[6] As we have said, the $r = 0$ case should vanish and $r = 1$ with a single $\langle Y \tau \rangle$ contraction should give the Einstein amplitudes at $\Lambda = 0$; $r > 1$ will give contributions to the de Sitter amplitudes that are higher order in Λ. An example of a Feynman tree with $r = 1$ contributing to the five-point MHV amplitude is given in figure 2.

We now investigate this formalism in detail for MHV amplitudes, and then generalize it to the N^{k}MHV case in a subsequent section.

---

[6] There are also potential powers of Λ coming from the Poisson structure, but as we observe later, these do not contribute at $O(1)$ or $O(\Lambda)$.



Figure 2. *An example of a Feynman tree diagram contributing to* $\mathcal{M}_5^0$

3.4 The $O(\Lambda^0)$ case

For $k = 0$ the diagrams will have 2 black and $n - 2$ white vertices. With $\Lambda = 0$ there will be no $\langle Y_i \tau_j \rangle$ contractions and the only edges in play are the directed ones from a vertex operator with a Y to any twistor wave function.

We begin by drawing a directed graph G which features *all* possible edges at once: each white vertex has $n - 1$ outgoing edges (see figure 3) to each of the other vertices, but just $n - 3$ incoming from the remaining white vertices. Each black vertex has just the incoming $n - 2$ edges from each of the white vertices. The $(n - 2)(n - 1)$ edges of G can be labeled by a pair (i, j) for the edge starting on white vertex i and ending on $j \neq i$.

The Matrix-Tree theorem in its simplest form allows us to count the number of oriented spanning trees of G; later we will weight the counting by the contribution from all the contractions. The key device is the $n \times n$ Laplacian matrix of G constructed as follows.

We first define the two modified incidence matrices, E and $\tilde{E}$ for G with rows indexed by the vertices v of G and columns by the edges e. The (v, e) entry of E is 1 if e is outgoing from v and 0 otherwise, whereas $\tilde{E}$ has $+1$ if e is outgoing and -1 if e is incoming and 0 otherwise. For G, the components are given by:

$$E_{i\ (j,k)} = \begin{cases} 1 \text{ if } i = j \\ 0 \text{ otherwise} \end{cases}, \qquad \tilde{E}_{i\ (j,k)} = \begin{cases} 1 \text{ if } i = j \\ -1 \text{ if } i = k \\ 0 \text{ otherwise} \end{cases}. \tag{3.2}$$

The Laplacian matrix is then defined by

$$L = \tilde{E}E^{\mathrm{T}} = \begin{pmatrix} n-1 & -1 & \cdots & -1 & 0 & 0 \\ -1 & \ddots & & \vdots & \vdots & \vdots \\ \vdots & & & n-1 & 0 & 0 \\ & & & -1 & 0 & 0 \\ -1 & \cdots & & -1 & 0 & 0 \end{pmatrix}. \tag{3.3}$$

The Matrix-Tree theorem (see appendix A or [23–25] for some background) enables us to use L to count the Feynman trees entering into the $O(\Lambda^0)$ part of the MHV correlator.

Lemma 3.1 *The number of non-trivial MHV Feynman tree diagrams at* $O(\Lambda^0)$ *is* $2 \det \tilde{L}_{n-1\, n} = 2(n-1)^{n-3}$, *where* $\tilde{L}_{n-1\, n}$ *is the matrix* L *with the* $n-1^{\text{th}}$ *and* n^{th} *rows and columns deleted, and* 1 *subtracted from the diagonal.*



Figure 3. *The graph G features all possible contractions within the $O(\Lambda^0)$ class*

Proof: The MHV Feynman graphs are spanning trees of the graph G obeying our rules (so that, in particular, there is only one outgoing edge from each white vertex). By Euler's theorem for a tree the $n-2$ edges connect $n-1$ vertices, so one vertex is left out, and it must be one of the black ones: either $n-1$ or n. Consider the class of graphs where n has no incoming edges: delete the n^{th} row and column from L and subtract one from the diagonal entries (as each white vertex no longer connects to vertex n) to obtain $\tilde{L}_n$. Now, let $\tilde{L}_n$ be the Laplacian matrix of some new graph G'. The Feynman graphs in this class are the spanning trees of G' rooted at vertex $n-1$; by the Matrix-Tree theorem (see appendix A), these are counted by $\det \tilde{L}_{n-1\,n}$. By Cayley's theorem, this is equal to $(n-1)^{n-3}$. The other class is counted in an equivalent fashion giving the same answer. □

To perform the sum of Feynman graphs in the correlator, we construct a weighted version $\mathcal{L}$ of L which includes the propagators required by the Feynman rules for the contractions. We insert a matrix over the set of edges whose diagonal entries are the twistor-string propagators:

$$\mathcal{P} = \text{diag}(\phi^i_j), \qquad \phi^i_j = \frac{[ij](\xi\, j)^2}{(ij)(\xi i)^2}, \tag{3.4}$$

so that we can define the weighted Laplacian as

$$\begin{aligned} \mathcal{L} &= \tilde{E}\mathcal{P}E^{\mathrm{T}}, \qquad\qquad (3.5)\\ &= \begin{pmatrix} \sum_{i\neq 1}\phi^1_i & -\phi^1_2 & \cdots & -\phi^1_{n-2} & 0 & 0 \\ -\phi^2_1 & \ddots & & \vdots & \vdots & \vdots \\ \vdots & & & \sum_{i\neq n-2}\phi^{n-2}_i & 0 & 0 \\ & & & -\phi^{n-1}_{n-2} & 0 & 0 \\ -\phi^n_1 & & \cdots & -\phi^n_{n-2} & 0 & 0 \end{pmatrix}. \end{aligned}$$

This differs from the matrix $\tilde{\phi}^0$ of (1.1) by a rank 2 error term, and from Hodges' matrix $\tilde{\Phi}$ (the $k=0$ form of (1.2)) by a conjugation and rank 2 error term:

$$\mathcal{L} = -\tilde{\phi}^0 + \mathcal{E} = -T^{-1}\tilde{\Phi}T + \mathcal{E}, \qquad \mathcal{E} = \begin{pmatrix} & -\phi^1_{n-1} & -\phi^1_n \\ \emptyset & \vdots & \vdots \\ & \sum_{i\neq n-1}\phi^i_{n-1} & -\phi^{n-1}_n \\ & -\phi^n_{n-1} & \sum_{i\neq n}\phi^i_n \end{pmatrix}, \qquad T = \text{diag}\left((\xi\, i)^2\right).$$

The matrix $\tilde{\Phi}$ has co-rank 3 thanks to the relation

$$\sum_{j=1}^{n} \tilde{\Phi}_{ij} \sigma_j^A \sigma_j^B = 0, \tag{3.6}$$

which follows from momentum conservation [13]. Furthermore, the error term is such that once row and columns $n-1$ and n are deleted, the matrices are conjugate. It is this high co-rank that will enable us to deduce that the sum of diagrams vanishes.

Although Lemma 3.1 generalizes to tell us how to compute the relevant correlators by taking determinants of minors of $\mathcal{L}$, we must divide up into sub-cases according to whether the remaining outgoing propagator from the white vertices lands on the black vertex $n-1$ or n. We first, consider the Feynman trees rooted at vertex $n-1$; this means that there can be no directed edges to vertex n and we must delete ϕ_n^i from each diagonal entry. We therefore define

Definition 3.1 *Let $\mathcal{L}^n$ be $\mathcal{L}$ with ϕ_n^i deleted from the ith diagonal entry. We further define $\mathcal{L}^n_{i \ldots k}$ to be $\mathcal{L}^n$ with rows and columns $i, \ldots, k$ deleted.*

For diagrams not involving the vertex n, we must also cross out the n^{th} row and column of $\mathcal{L}$ leaving the modified $(n-1) \times (n-1)$ matrix denoted $\mathcal{L}^n_n$. The sum of correlators associated to diagrams in this class is then given by crossing out the row and column of $\mathcal{L}^n_n$ corresponding to vertex $n-1$ to yield $\mathcal{L}^n_{n-1\, n}$ and taking its determinant, thanks to the weighted version of the Matrix-Tree theorem (c.f., appendix A or [24], Chapter 36). A similar pattern follows for the other class of trees rooted at vertex n. This indicates that the $\Lambda = 0$ portion of the conformal gravity MHV correlator is given by

$$\mathcal{C}_{n,1}|_{O(\Lambda^0)} = \det \mathcal{L}^n_{n\, n-1} + \det \mathcal{L}^{n-1}_{n-1\, n}.$$

This sum can be re-interpreted thanks to some basic facts about the matrix $\mathcal{L}$ and determinants.

Lemma 3.2 *We have*

$$\det \mathcal{L}^{\cdots i}_{\ldots i\, j} = \sum_{k \neq \ldots i, j} R_k \phi_j^k\,, \qquad i, j, k = 1, \ldots, n\,, \tag{3.7}$$

where $R_k = \mathcal{L}^{\cdots i}_{\ldots ijk}$.

Proof: The matrix $\mathcal{L}^{\cdots i}_{\ldots ij}$ only depends on the ϕ_j^k through its diagonal entries by the definition of $\mathcal{L}$ in (3.5). We can therefore expand its determinant out along the diagonal by

$$\det \mathcal{L}^{\cdots i}_{\ldots i\, j} = A + \sum_{k \neq i, j} R_k \phi_j^k,$$

where $R_k = \mathcal{L}^{\cdots i}_{\ldots ijk}$ so it suffices to show that $A = 0$. By lemma 3.1 and the Matrix-Tree theorem, we know that each term in $\det \mathcal{L}^{\cdots i}_{\ldots i\, j}$ corresponds to the CFT correlator of a tree rooted at vertex j (omitting vertices … also). This means that every term will contain (at least) one propagator to vertex j, by definition of a rooted tree graph. Thus there can be no term in $\det \mathcal{L}^{\cdots i}_{\ldots i\, j}$ of order zero in ϕ_j^k for all $k \neq j$; in other words, $A = 0$. $\Box$

Lemma 3.3 $\det \mathcal{L}^{n}_{n\,n-1} + \det \mathcal{L}^{n-1}_{n-1\,n} = \det \mathcal{L}_{n-1\,n}$.

Proof: Lemma 3.2 means that,

$$\det \mathcal{L}^{n}_{n\,n-1} = \sum_{i\neq n-1,n} R_i \phi^i_{n-1}, \qquad \mathcal{L}^{n-1}_{n-1\,n} = \sum_{i\neq n-1,n} S_i \phi^i_{n},$$

where R_i and S_i are the remaining factors. However, the matrices $\mathcal{L}^{n}_{n\,n-1}$ and $\mathcal{L}^{n-1}_{n-1\,n}$ differ only by the presence of ϕ^i_{n-1} and ϕ^i_n along their diagonals, respectively, so it follows that $R_i = S_i$. Therefore, we can write

$$\det \mathcal{L}^{n}_{n\,n-1} + \det \mathcal{L}^{n-1}_{n-1\,n} = \sum_{i\neq n-1,n} R_i (\phi^i_{n-1} + \phi^i_n) = \det \mathcal{L}_{n-1\,n},$$

as required. □

Lemmas 3.2 and 3.3 therefore give

$$\det \mathcal{L}_{n-1\,n} = \det \left(-(T^{-1}\tilde{\Phi}T)_{n-1\,n} + \mathcal{E}_{n-1\,n} \right) = (-1)^{n-2} \det \tilde{\Phi}_{n-1\,n} = 0, \tag{3.8}$$

since $\tilde{\Phi}$ has co-rank three and the error term was only non-vanishing in the $n-1^{\text{st}}$ and n^{th} columns. This confirms that the $\Lambda = 0$ contribution to $\mathcal{C}_{n,1}$ vanishes, as required by the Maldacena relation between conformal and Einstein gravity amplitudes.

4 The MHV Amplitude From the $O(\Lambda)$ Contribution

Hodges' formula for the MHV amplitude in $\mathcal{N} = 8$ supergravity is [13]:

$$\mathcal{M}^0_n = \delta^{4|16}\left(\sum_{j=1}^{n} \lambda_j \tilde{\lambda}_j \right) \det{}' \left(\tilde{\Phi}^0 \right), \tag{4.1}$$

where $\det'(\tilde{\Phi}^0)$ is a reduced determinant of the MHV ($k = 0$) matrix (1.2), and the delta-function expresses super-momentum conservation. Although other forms of the MHV amplitude have been known for some time (e.g., the BGK or BCFW type formulas [27–29]), (4.1) manifests the full permutation symmetry of gravitational scattering amplitudes without a cumbersome summation, and makes no reference to any cyclic ordering of the gravitons. In this sense, Hodges' formula is the closest gravitational analogue to the Parke-Taylor formula for gluon scattering in Yang-Mills theory. It was proved using a $\mathcal{N} = 7$ version of BCFW recursion [30]. Here we will see how it arises from the twistor-string formula (2.9) at $O(\Lambda)$.

There are two possible $O(\Lambda)$ contributions. The first arises from a Y-τ contraction as we described in the previous section. The other arises from the $O(\Lambda)$ part of the Poisson structure and will potentially contribute to all possible contractions between a Y and a wavefunction. The second can be treated very similarly to the $O(\Lambda^0)$ case and can be seen to have a vanishing contribution.



Figure 4. *The graph G^i showing all possible contractions subject to a given $\langle Y_i \tau_n \rangle$ contraction*

We first of all ignore the τs so the contribution from the downstairs infinity twistor is irrelevant (the vanishing will be seen before any integration of a τ is performed as in the previous section). The argument for the vanishing of the $O(\Lambda^0)$ contribution to the correlator works for any choice of wave functions, so the calculation is essentially conformally invariant (we are working in a conformal gravity framework) and will work to show that the contribution of the connected tree diagrams with no Y-τ contractions vanishes for any choice of simple upstairs infinity twistor. Thus the derivative of the correlator with respect to the infinity twistor vanishes in any direction for which the infinity twistor remains simple. But such directions span all possible variations in the infinity twistor: an infinity twistor corresponds a point in $\mathbb{C}^6$ and the simple ones to lightcones in $\mathbb{C}^6$ and any direction can be expressed as a sum of null directions. In particular, the derivative with respect to Λ must vanish and so the $O(\Lambda)$ term vanishes.[7]

Hence, the $O(\Lambda)$ contribution to the correlator is generated by those Feynman trees which have a single $\langle Y_i \tau_j \rangle$ contraction. Without loss of generality, suppose this contraction occurs between vertex i and n. We will see that, remarkably, once the answer is added on to the case where Y_i contracts with τ_{n-1} the result is independent of the choice of i.

We proceed to draw a graph G^i that contains all the remaining possible edges allowed by our Feynman rules (see figure 4). From this point, we can proceed almost identically to the $O(\Lambda^0)$ case. The modified incidence matrices now read

$$E^i_{j\ (k,l)} = \begin{cases} 1 \text{ if } j = k \text{ and } j \neq i \\ 0 \text{ otherwise} \end{cases}, \qquad \tilde{E}^i_{j\ (k,l)} = \begin{cases} 1 \text{ if } j = k \text{ and } j \neq i \\ -1 \text{ if } j = l \\ 0 \text{ otherwise} \end{cases}, \tag{4.2}$$

and the weighted Laplacian matrix $\mathcal{L} = \tilde{E}^i \mathcal{P} (E^i)^{\mathrm{T}}$ is the same as in (3.5) except that now the i^{th} column has all zero entries.

The contraction $\langle Y_i \tau_n \rangle$ combined with $\langle Y_i \tau_{n-1} \rangle$ produces a factor corresponding to the 3-point MHV amplitude. We will discuss this factor in greater detail later, but we will take its value[8] (with $\mathcal{N}=4$ supersymmetry) to be given by:

---

[7] An alternative argument using dual-twistor wave functions can be used to show directly that any choice of upstairs infinity twistor gives zero when Y-τ contractions are ignored using precisely analogous arguments as the previous subsection but now with a Hodges like matrix in which the numerators are replaced by $I^{\alpha\beta} W_{i\alpha} W_{j\beta}$ and the co-rank 3 arises from an analogue of momentum conservation arising from the delta-functions in the Grassmannian approach of [6].

$$\langle Y_i \tau_n \tau_{n-1}\rangle = \Lambda \mathrm{D}\sigma_i \mathrm{D}\sigma_n \mathrm{D}\sigma_{n-1} \frac{(n\ n-1)^2}{(i\ n)^2 (n-1\ i)^2}. \tag{4.3}$$

Hence, we are left with the task of computing the factor arising from the remaining sum over trees in G^i.

There are $n-3$ remaining Y_is and these must connect the remaining $n-3$ vertices into a tree so there will be one remaining outgoing contraction from this tree and this can contract with the vertex i, $n-1$ or n. The arguments in the proof of lemma 3.1 show that there will be $(n-2)^{n-4}$ trees contributing, and that the sum of correlators is given by:

$$\det \mathcal{L}^{i\ n-1}_{i\ n-1\ n} + \det \mathcal{L}^{n-1\ n}_{n-1\ n\ i} + \det \mathcal{L}^{n\ i}_{n\ i\ n-1}, \tag{4.4}$$

Here, $\mathcal{L}^{i\,j}_{i\,j\,k}$ is the matrix $\mathcal{L}$ with the i^{th}, j^{th}, and k^{th} rows and columns deleted, and the ϕ^i_l, ϕ^j_l terms subtracted from the remaining diagonal entries. A simple application of lemma 3.2 tells us that

$$\det \mathcal{L}^{i\ n-1}_{i\ n-1\ n} = \sum_j R_j \phi^j_n, \quad \det \mathcal{L}^{n-1\ n}_{n-1\ n\ i} = \sum_j R_j \phi^j_i, \quad \det \mathcal{L}^{n\ i}_{n\ i\ n-1} = \sum_j R_j \phi^j_{n-1},$$

and by lemma 3.3, this gives

$$\det \mathcal{L}^{i\ n-1}_{i\ n-1\ n} + \det \mathcal{L}^{n-1\ n}_{n-1\ n\ i} + \det \mathcal{L}^{n\ i}_{n\ i\ n-1} = \sum_j R_j (\phi^j_i + \phi^j_{n-1} + \phi^j_n) = \det \mathcal{L}_{i\ n-1\ n}.$$

Noting that $\det \mathcal{L}^i_{i\ n-1\ n} = (-1)^{n-3} \det \tilde{\Phi}_{i\ n-1\ n}$, we then find that the correlator with a single contraction of the form $\langle Y_i \tau_n \rangle$ gives a contribution:

$$\mathcal{C}^i_{n,1}|_{O(\Lambda)} = \Lambda \frac{(n\ n-1)^2}{(i\ n)^2 (n-1\ i)^2} \det \tilde{\Phi}_{i\ n-1\ n}. \tag{4.5}$$

Although we still have to sum over the $n-2$ different choices of i for the $\langle Y_i \tau_n \rangle$ contraction (the interchange $n \leftrightarrow n-1$ is already taken care of in (4.3)), the defining properties of the reduced determinant $\det'$ of the matrix $\tilde{\Phi}$ in fact show that all of these terms will be equal [13]. This is a consequence of the emergent permutation symmetry that arises in $\mathcal{N}=8$ supergravity.

Indeed, we can introduce a power of $(n-1\ n)^{-4}$ into the three point MHV amplitude (4.3) by extending the supersymmetry to $\mathcal{N}=8$ where now this emergent permutation symmetry will extend over n and $n-1$ also. This gives the $\mathcal{N}=8$ momentum-space amplitude:

$$\begin{aligned}\mathcal{M}^0_n = \int_{\mathscr{M}_{1,n}} \frac{\mathrm{d}^{4|8}U_0 \wedge \mathrm{d}^{4|8}U_1}{\mathrm{vol\ GL}(2,\mathbb{C})} \prod_{j=1}^n \mathrm{D}\sigma_j h_j(\lambda_j \tilde{\lambda}_j; Z(\sigma_j)) \frac{\det \tilde{\Phi}_{i\ n-1\ n}}{(n\ n-1)^2 (n-1\ i)^2 (i\ n)^2} \\ = \delta^{4|16}\left(\sum_{j=1}^n \lambda_j \tilde{\lambda}_j\right) \det{}'(\tilde{\Phi}^0).\end{aligned} \tag{4.6}$$

---

[8] We will see later that our naïve rules that we have used here give rise to a gauge-dependent error term.

Here, the super-momentum-conserving delta function is realized via Nair's lemma [31]

$$\delta^{4|16}\left(\lambda\tilde{\lambda}\right) = \int \mathrm{d}^{4|16}x \, \exp\left(i\lambda_A\tilde{\lambda}_{A'}x^{AA'} + i\eta_a\lambda_A\theta^{Aa}\right),$$

and $\det'(\tilde{\Phi}^0)$ is short-hand for:

$$\det{}'(\tilde{\Phi}^0) \equiv \frac{\det(\tilde{\Phi}_{i\,n-1\,n})}{\langle i\; n-1\rangle^2\langle n-1\; n\rangle^2\langle n\; i\rangle^2},$$

with the $\mathrm{GL}(2,\mathbb{C})$ freedom fixed by setting $\sigma_j = \lambda_j$.

Equation (4.6) is just a particular representation of Hodges' formula (4.1) in the form found by Cachazo and Skinner [5, 6], so our Feynman tree formalism correctly produces the MHV amplitude for gravity.

4.1 The 3-point amplitude

In arriving at (4.6), we took for granted that our twistor-string theory correctly reproduced the three-point MHV amplitude with $\mathcal{N} = 4$ supersymmetry:

$$\mathcal{M}_3^0 = \frac{\langle n-1\; n\rangle^2}{\langle i\; n-1\rangle^2\langle n\; i\rangle^2}, \tag{4.7}$$

where we have suppressed a momentum conserving delta-function. The Feynman diagram associated with the three point amplitude is depicted in figure 5, and the twistor-string expression for the correlator is given by:

$$\int_{\mathscr{M}_{1,3}} \mathrm{d}\mu_1 \left\langle Y_i \cdot \partial h_i \; \tilde{h}_{n-1}\tau_{n-1} \; \tilde{h}_n\tau_n \right\rangle. \tag{4.8}$$

However, the naïve use of Wick contractions in the worldsheet CFT leads to a gauge dependent error term between (4.3) and (4.8) given by

$$\frac{\langle\xi 1\rangle\langle\xi 3\rangle}{\langle 12\rangle\langle 23\rangle\langle\xi 2\rangle^2}. \tag{4.9}$$

This same error arises also if we directly use the three point formula given by Nair [18].

In [9] we instead used a geometric computation based on a technique utilized to construct the MHV amplitude in the Einstein case in [28]. This was based on the conformal gravity twistor action of [14] which closely models the twistor-string theory calculation. It is clear that the naïve twistor-string rules that we have been using are not quite correct in this context given that they are leading to a gauge-dependent answer.

In summary, if we assume that twistor-string theory produces the correct three-point MHV amplitude, then we have a complete derivation of Hodges' MHV formula from twistor-string theory Feynman rules. However, there is a gap in our understanding of the three-point correlator in the world-sheet CFT and in particular the $\langle Y\,\tau\rangle$ contractions which are not yet properly understood.



Figure 5. *One of the two graphs of the three-point amplitude* $\langle Y_i \cdot \partial h_i \tilde{h}_n \tau_n \tilde{h}_{n-1} \tau_{n-1} \rangle$

5 The N^{k}MHV Amplitude

At N^{k}MHV, the Hodges MHV formula is generalised by the Cachazo-Skinner formula [5, 6]. Like the twistor-string formula, it is an integral over the moduli of rational curves of degree $d = k - 1$. It provides all tree-level N^{k}MHV amplitudes of $\mathcal{N} = 8$ supergravity [6] and is given by:

$$\mathcal{M}_n^k(1, \ldots, n) = \int \frac{\prod_{r=0}^{k+1} \mathrm{d}^{4|8} U_r}{\text{vol GL}(2, \mathbb{C})} \det{}'(\tilde{\Phi}^k) \det{}'(\Phi^k) \prod_{i=1}^{n} \mathrm{D}\sigma_i \; h_i \left(\lambda_i \tilde{\lambda}_i ; Z(\sigma_i) \right) . \tag{5.1}$$

Here, $\tilde{\Phi}^k$ is the co-rank $k + 3$ matrix (1.2), and Φ^k is essentially its parity conjugate:

$$\Phi_{ij} = \begin{cases} \frac{\langle ij \rangle}{(i\, j)} & i \neq j \\ -\sum_{l \neq i} \frac{[i\, l]}{(i\, l)} \left(\frac{(\xi l)}{(\xi i)} \right)^{n-k-1} \frac{\prod_{m \neq i} (i\, m)}{\prod_{s \neq l} (l\, s)} & i = j \end{cases} , \tag{5.2}$$

The $\det'(\tilde{\Phi}^k)$ and $\det'(\Phi^k)$ are modified reduced determinants, involving some additional Vandermonde determinant factors. In particular, $\det'(\tilde{\Phi}^k)$ can be written

$$\det{}'(\tilde{\Phi}^k) = \frac{\det \left(\tilde{\Phi}^k_{i\, n-k-2 \cdots n} \right)}{\prod_{j<k,\; j,k \in \{i, n-k-2, \ldots, n\}} (j\; k)} \equiv \frac{\det \left(\tilde{\Phi}^k_{i\, n-k-2 \cdots n} \right)}{\mathrm{VM}(i, n-k-2, \ldots, n)} . \tag{5.3}$$

That (5.1) involves an integral over the moduli space of n-pointed stable maps of $\mathbb{CP}^1$ to twistor space is of course highly suggestive of a twistor-string origin. We will here not attempt to explain the whole formula, but just the factor $\det \tilde{\Phi}^k_{i\, n-k-2 \cdots n}$ in terms of our tree graph arguments generalized from the MHV setting. The remaining terms should presumably also arise from the other ingredients in the twistor-string calculations but we have not yet successfully identified those terms.

For the N^{k}MHV amplitude the twistor-string tree formula (2.9)

$$\lim_{\Lambda \to 0} \frac{1}{\Lambda} \int_{\mathscr{M}_{k+1,n}} \mathrm{d}\mu_{k+1} \; \mathcal{C}_{n,k+1} .$$

is now an integral over the moduli of rational maps of degree $d = k - 1$ and there are now $n - k - 2$ vertex operators of the form $Y \cdot \partial h$ and $k + 2$ of the form $\tilde{h}\tau$. Again we will allow only maximally connected Feynman tree diagrams to contribute to the CFT correlator $\mathcal{C}_{n,k+1}$; if we do not, then we will again see that we obtain incorrect non-vanishing answers at $\Lambda = 0$. The diagrams therefore now have $k + 2$ black vertices and $n - k - 2$ white vertices. In particular there are just $n-k-2$ Y_is to be contracted with the h_j, $\tilde{h}_j$ or τ_j and so the diagrams become correspondingly simpler with fewer edges. The number of edges corresponds to the size of the minors whose determinant we must take when we use the Matrix-Tree theorem. It turns out that the ranks of the matrices $\tilde{\Phi}^k$ given in (1.1) and (1.2) encoding the twistor-string propagators are $n-k-3$, decreasing as appropriate with k.

Again at $\Lambda = 0$, where we must have vanishing contribution, there can be no $\langle Y \tau \rangle$ contractions and for the $O(\Lambda)$ contribution (which should give the Einstein amplitude) there should be just one. As before, we draw the master graph G of all possible Y_i contractions with the h_j and $\tilde{h}_j$. Thus in G, black vertices have $n-k-2$ incoming edges (one from each of the white vertices) and no outgoing edges, while white vertices will have $n-k-3$ incoming (one from each of the other white vertices) and $n-1$ outgoing edges to all other vertices. The definition of the modified incidence matrices E, $\tilde{E}$ and $L = \tilde{E}E^{\mathrm{T}}$ for this new graph follows in precisely the same way as in the MHV setting.

The incidence Laplace matrix of G is now:

$$L = \begin{pmatrix} n-1 & -1 & \cdots & -1 & 0 & \cdots & 0 \\ -1 & \ddots & & \vdots & \vdots & & \vdots \\ \vdots & & & n-1 & 0 & \cdots & 0 \\ & & & -1 & 0 & \cdots & 0 \\ & & & \vdots & \vdots & & \vdots \\ -1 & \cdots & & -1 & 0 & \cdots & 0 \end{pmatrix} ,$$

with the last $k+2$ columns composed entirely of zeros. For the propagators to form a connected tree, there can only be one outgoing propagator from the white vertices. If this lands on vertex $n-k-2$, the other black vertices are isolated and the Matrix-Tree theorem gives the number of such Feynman trees rooted at the black vertex $n-k-2$ to be $\det \tilde{L}_{n-k-2\cdots n} = (n-k-1)^{n-k-3}$.

To compute the actual correlator at $O(\Lambda^0)$ rather than just the number of contributing trees, we define the weighted Laplacian $\mathcal{L}$ weighted by the appropriate propagators at degree $d = k+1$ which gives

$$\mathcal{L} = \begin{pmatrix} \sum_{i\neq 1} \phi^1_i & -\phi^1_2 & \cdots & -\phi^1_{n-k-2} & 0 & \cdots & 0 \\ -\phi^2_1 & \ddots & & \vdots & \vdots & & \vdots \\ \vdots & & & \sum_{i\neq n-k-2} \phi^{n-k-2}_i & 0 & \cdots & 0 \\ & & & -\phi^{n-k-1}_{n-k-2} & \vdots & & \vdots \\ & & & \vdots & & & \\ -\phi^n_1 & \cdots & & -\phi^n_{n-k-2} & 0 & \cdots & 0 \end{pmatrix} , \qquad \phi^i_j = \frac{[i\,j]}{(i\,j)} \left(\frac{(\xi\,j)}{(\xi\,i)} \right)^{k+2} .$$

This $\mathcal{L}$ is related to $\tilde{\Phi}^k$ by a conjugation by $T = \mathrm{diag}\left((\xi\,i)^{k+2} \right)$ and a rank $k+2$ error term that disappears when the last $n-k-2$ rows and columns are deleted. The natural generalizations of lemmas 3.2 and 3.3 then tell us that the sum of the Feynman trees contributing to the correlator $\mathcal{C}_{n,k+1}$ is given by:

$$\det \mathcal{L}_{n-k-2\cdots n} = \det \tilde{\Phi}^k_{n-k-2\cdots n} = 0,$$

and this is vanishing because $\tilde{\Phi}^k$ has co-rank $k+3$. This gives the expected vanishing at $\Lambda = 0$.

The Einstein gravity contribution should be given by the $O(\Lambda)$ contribution where there is just a single $\langle Y_i \tau_j \rangle$ contraction, as in the MHV case. Instead of a 3-point MHV pre-factor, we expect to obtain something more complicated which encodes the data of the single contraction $\langle Y_i \tau_j \rangle$, the τs and other Jacobian fators[9]. We will ignore such factors here and focus only from those that arise from contractions of the remaining Ys with the hs and $\tilde{h}$s.

As in the MHV amplitude, this amounts to trees rooted at one of vertex i or any of the $k+2$ τ-vertices. Yet again, counting these $(n-k-2)^{n-k-4}$ trees corresponds to taking reduced determinants of the matrix $\mathcal{L}$ via the natural extensions of lemmas 3.1-3.3. Hence, we obtain the contribution $\det \mathcal{L}_{i\, n-k-2\cdots n} = \det \tilde{\Phi}^k_{i\, n-k-2\cdots n}$. Of course, we must still sum over all of the different ways of taking the initial $\langle Y_i \tau_j \rangle$ contribution, but as in the MHV case, the properties of the $\tilde{\Phi}^k$ matrix ensure that all of these contributions are equal up to the kind of Vandermonde factors that we are ignoring.

6 Discussion

Although our derivation is far from complete, we have nevertheless made significant progress in our check that the Berkovits-Witten twistor-string correctly reproduces the Einstein subsector of conformal gravity amplitudes. Now that we have the Cachazo-Skinner formula (5.1), our task has been to identify how terms in that formula can arise from twistor-string theory. We clearly have an explicit derivation of not only the moduli space integral and the external wave functions, but also the reduced determinant of $\tilde{\Phi}^k$ in the formula. This arose by requiring that only connected tree diagrams should be taken to contribute to the CFT worldsheet correlators arising in the twistor-string amplitude formula; the reduced determinant arose by expressing the sum of the resulting Feynman diagrams as the determinant of a weighted Laplacian matrix for the graph of all possible contractions using the weighted Matrix-Tree theorem. Although this provides substantial, perhaps overwhelming, circumstantial evidence that the Berkovits-Witten twistor-string correctly gives the (tree-level) gravity sub-sector of conformal gravity, and hence circumstantial evidence that all of conformal gravity can be obtained in this way, we are still missing complete arguments even for a part (equation (4.9)) of the Hodges gravitational MHV formula. At this stage our argument does not reproduce any of $\det' \Phi$ nor the Vandermonde determinants in the N^{k}MHV Cachazo-Skinner formula.

At this stage it seems likely that the Vandermonde determinants and $\det' \Phi$ factors must all be present in the Berkovits-Witten twistor-string formula. However, a better understanding of the worldsheet conformal field theory is required even for the three point amplitude.

---

[9] In the $k=0$, or $d=1$ case, we were free to fix the GL$(2,\mathbb{C})$ freedom so that $\sigma_i = \lambda_i$; thereby setting the Jacobian factor equal to unity. For $d>1$ this is no longer possible and a non-trivial factors can occur.

It is somewhat vexing that only tree diagrams are required in the worldsheet correlator. From the string theory perspective this is difficult to motivate as there are no parameters that can be used to suppress loop contributions. There may be some exponentiation argument as in the Yang-Mills case [32]: there the twistor-string naïvely gives the determinant of the twistor $\bar{\partial}$-operator whereas the correct contribution to obtain Yang-Mills is the log of the determinant. However, the geometric interpretation of this restriction to trees is clear. The Y contractions are performing a perturbation expansion for the construction of holomorphic curves in a deformed twistor space given as solutions to

$$\bar{\partial}_\sigma Z^\alpha(x,\sigma) = I^{\alpha\beta}\partial_\beta h(Z)\,. \tag{6.1}$$

The Feynman trees build up the correct classical solution to this equation as appropriate to the Penrose non-linear graviton construction. The loop contributions would be appropriate to some quantisation of this construction, whereas all we are trying to obtain here are classical solutions in order to get gravity tree amplitudes (quantum ones according to string theory should only arise when higher genus curves are introduced).

We have presented this work as a verification of the Berkovits-Witten twistor-string, but it is also the case that a key motivation is to elucidate the twistor-string underpinnings of the Cachazo-Skinner formula. Although it is very suggestive of a twistor-string theory that directly describes $\mathcal{N}=8$ supergravity, the details are far from clear. The ingredients described in this paper will be very likely to play a role also in such a theory, or at least in its $\mathcal{N}=4$ truncation, which should in turn embed in the conformal gravity twistor-string.

Acknowledgments

We would like to thank Mat Bullimore, Freddy Cachazo, Song He, Andrew Hodges, Parameswaran Nair and David Skinner for useful conversations and comments. While completing this project, we heard about related work in progress by Bo Feng and Song He [33] and Clifford Cheung [34] relating the MHV tree formula of [12] to Hodges' formula. TA is supported by a National Science Foundation (USA) Graduate Research Fellowship and by Balliol College; LM is supported by a Leverhulme Fellowship and benefited from hospitality in the Perimeter Institute during some of the work on this project.

A Graph Theory

In this appendix, we review some graph theory that we have used to sum the Feynman tree graphs. For a more complete introduction see e.g., [23–25].

Let G be a directed graph, with a set of vertices $\mathcal{V}$ and directed edges $\mathcal{E}$. We will denote such a graph by $G=(\mathcal{V},\mathcal{E})$. Denote the number of vertices as $|\mathcal{V}|=n$ and edges as $|\mathcal{E}|=m$. The *incidence matrix* of G is a $n\times m$ matrix E encoding the connective structure of the graph. We can denote an edge of G as $(j,k)\in\mathcal{E}$, meaning the directed edge which is outgoing from vertex $j \in \mathcal{V}$ and incoming to vertex $k \in \mathcal{V}$. In this notation, the entries of the incidence matrix are given by:

$$E_{i\ (j,k)} = \begin{cases} \ \ 1 \text{ if } i = j \\ -1 \text{ if } i = k \\ 0 \text{ otherwise} \end{cases} . \tag{A.1}$$

The incidence matrix can be used to build the *Laplacian matrix* of the graph G, which is a $n \times n$ matrix given by:

$$\Delta = E E^{\mathrm{T}}, \qquad \Delta_{i\,j} = \begin{cases} -1 \text{ if } i \neq j \text{ and } (i,j) \in \mathcal{E} \\ \quad d_i^{\text{out}} \text{ if } i = j \\ \quad 0 \text{ otherwise} \end{cases} , \tag{A.2}$$

where d_i^{out} is the out-degree of vertex $i \in \mathcal{V}$.

The Feynman tree diagrams we are interested in counting throughout this paper are examples of particular sub-graphs of a directed graph, namely rooted spanning trees:

Definition A.1 *A* spanning tree of G rooted at $v \in \mathcal{V}$ *is a spanning sub-graph $T = (\mathcal{V}, \mathcal{F})$ of G such that: (1.) T has no oriented cycles; (2.) vertex v has out-degree zero; and (3.) every other vertex $i \neq v$ has out degree one.*

The main result in graph theory we use is the weighted generalisation of the Matrix-Tree theorem. The original matrix-tree theorem tells us how to count spanning trees of G rooted at a particular vertex in terms of determinants of the reduced Laplacian of G (c.f., [25], Chapter 9; [24], Chapter 36). Since it illustrates the enumerative nature of the arguments used in the body of the text, we provide a proof here:

Theorem 1 (Matrix-Tree Theorem) *Let $\kappa(G, v)$ denote the number of oriented spanning trees of G rooted at $v \in \mathcal{V}$, and Δ_v be the Laplacian matrix of G with the v^{th} row and column deleted. Then $\kappa(G, v) = \det \Delta_v$.*

Proof: Without loss of generality, let $v = n \in \mathcal{V}$. Let the $(n-1) \times n$ matrix F be the incidence matrix E of G with its n^{th} row deleted, so that $\Delta_n = F F^{\mathrm{T}}$. Now, the Cauchy-Binet theorem (c.f., [25], Chapter 9) tells us that we can write:

$$\det \Delta_n = \sum_S \det(F_S) \det(F_S^{\mathrm{T}}) = \sum_S \det(F_S)^2,$$

where S ranges over all subsets of $\mathcal{E}$ of size $n-1$ none of which are outgoing from $n \in \mathcal{V}$, and F_S is the $(n-1) \times (n-1)$ matrix whose columns are those of F with indices in S. This means that each choice of S corresponds of a sub-graph $T \subset G$ with $n-1$ edges which have $d_n^{\text{out}} = 0$. We now quote the following lemma from graph theory:

Lemma A.1 (c.f., [25] Lemma 9.7) *Let S be a set of $n-1$ directed edges of $G = (\mathcal{V}, \mathcal{E})$, none of which are outgoing from $n \in \mathcal{V}$. Then $\det F_S = \pm 1$ if S forms a spanning tree of G rooted at n, and $\det F_S = 0$ otherwise.*

This means that

$$\det \Delta_n = \sum_{\text{spanning trees rooted at n}} (\pm 1)^2 = \kappa(G,n),$$

as required. □

The weighted generalizations of the Matrix-Tree theorem include weighted edges (e.g., [24], Chapter 36) and allow us to introduce the propagator appropriate to each edge. This leads to the weighted Laplacians used throughout the paper. We define the weighted Laplacian matrix $\widehat{\Delta}$ for the directed graph G with weights for each edge given by:

$$\widehat{\Delta} = E\mathcal{W}E^{\mathrm{T}}, \mathcal{W} = \operatorname{diag}(w_{ij}), \widehat{\Delta}_{ij} = \begin{cases} -w_{ij} \text{ if } i \neq j \text{ and } (i,j) \in \mathcal{E} \\ \sum_{(i,k)\in\mathcal{E}} w_{ik} \text{ if } i = j \\ 0 \text{ otherwise} \end{cases}.$$

Then the weighted Matrix-Tree theorem reads:

Theorem 2 (Weighted Matrix-Tree Theorem) *Let $\mathcal{T}(G,v)$ be the set of all oriented spanning trees of G rooted at $v \in \mathcal{V}$ and $\widehat{\Delta}$ be the weighted Laplacian of G. For any $T \in \mathcal{T}(G,v)$, let $\mathcal{E}_T$ denote its set of directed edges. Then*

$$\det \widehat{\Delta}_v = \sum_{T\in\mathcal{T}(G,v)} \left(\prod_{(i,j)\in\mathcal{E}_T} w_{ij} \right).$$

When the weights are set to be twistor-string propagators, this is the primary result used in the text.

References

[1] E. Witten, *Perturbative gauge theory as a string theory in twistor space*, *Commun.Math.Phys.* **252** (2004) 189–258, [`hep-th/0312171`].

[2] R. Roiban, M. Spradlin, and A. Volovich, *On the tree level S matrix of Yang-Mills theory*, *Phys.Rev.* **D70** (2004) 026009, [`hep-th/0403190`].

[3] T. Adamo, M. Bullimore, L. Mason, and D. Skinner, *Scattering Amplitudes and Wilson Loops in Twistor Space*, *J.Phys.A* **A44** (2011) 454008, [`arXiv:1104.2890`].

[4] A. Brandhuber, B. Spence, and G. Travaglini, *Tree-Level Formalism*, *J.Phys.A* **A44** (2011) 454002, [`arXiv:1103.3477`].

[5] F. Cachazo and D. Skinner, *Gravity from Rational Curves*, `arXiv:1207.0741`.

[6] F. Cachazo, L. Mason, and D. Skinner, *Gravity in Twistor Space and its Grassmannian Formulation*, `arXiv:1207.4712`.

[7] M. Bullimore, *New Formulae for Gravity Amplitudes: Parity Invariance and Soft Limits*, `arXiv:1207.3940`.

[8] S. He, *A Link Representation for Gravity Amplitudes*, `arXiv:1207.4064`.

[9] T. Adamo and L. Mason, *Einstein supergravity amplitudes from twistor-string theory*, *Class.Quant.Grav.* **29** (2012) 145010, [`arXiv:1203.1026`].

[10] J. Maldacena, *Einstein Gravity from Conformal Gravity*, `arXiv:1105.5632`.

[11] N. Berkovits and E. Witten, *Conformal supergravity in twistor-string theory*, *JHEP* **0408** (2004) 009, [`hep-th/0406051`].

[12] D. Nguyen, M. Spradlin, A. Volovich, and C. Wen, *The Tree Formula for MHV Graviton Amplitudes*, *JHEP* **1007** (2010) 045, [`arXiv:0907.2276`].

[13] A. Hodges, *A simple formula for gravitational MHV amplitudes*, `arXiv:1204.1930`.

[14] L. Mason, *Twistor actions for non-self-dual fields: a new foundation for twistor-string theory*, *JHEP* **0510** (2005) 009, [`hep-th/0507269`].

[15] L. Mason and D. Skinner, *Heterotic twistor-string theory*, *Nucl.Phys.* **B795** (2008) 105–137, [`arXiv:0708.2276`].

[16] N. Berkovits, *An Alternative string theory in twistor space for N=4 superYang-Mills*, *Phys.Rev.Lett.* **93** (2004) 011601, [`hep-th/0402045`].

[17] T. Adamo and M. Groechenig, *Moduli stacks of maps for supermanifolds*, `arXiv:1204.4369`.

[18] V. Nair, *A Note on graviton amplitudes for new twistor string theories*, *Phys.Rev.* **D78** (2008) 041501, [`arXiv:0710.4961`].

[19] C.-h. Ahn, *Comments on MHV tree amplitudes for conformal supergravitons from topological B-model*, *JHEP* **0507** (2005) 004, [`hep-th/0504109`].

[20] L. Dolan and J. N. Ihry, *Conformal Supergravity Tree Amplitudes from Open Twistor String Theory*, *Nucl.Phys.* **B819** (2009) 375–399, [`arXiv:0811.1341`].

[21] M. Wolf, *Self-Dual Supergravity and Twistor Theory*, *Class.Quant.Grav.* **24** (2007) 6287–6328, [`arXiv:0705.1422`].

[22] L. Mason and M. Wolf, *Twistor Actions for Self-Dual Supergravities*, *Commun.Math.Phys.* **288** (2009) 97–123, [`arXiv:0706.1941`].

[23] R. P. Stanley, *Enumerative Combinatorics*, vol. 2. Cambridge University Press, 1999.

[24] J. van Lint and R. Wilson, *A Course in Combinatorics*, vol. 2. Cambridge University Press, 2001.

[25] R. P. Stanley, “Topics in Algebraic Combinatorics.” http://www-math.mit.edu/ rstan/algcomb.pdf, 2012.

[26] Z. Bern, L. J. Dixon, M. Perelstein, and J. Rozowsky, *Multileg one loop gravity amplitudes from gauge theory*, *Nucl.Phys.* **B546** (1999) 423–479, [`hep-th/9811140`].

[27] F. A. Berends, W. Giele, and H. Kuijf, *On relations between multi - gluon and multigraviton scattering*, *Phys.Lett.* **B211** (1988) 91.

[28] L. Mason and D. Skinner, *Gravity, Twistors and the MHV Formalism*, *Commun.Math.Phys.* **294** (2010) 827–862, [`arXiv:0808.3907`].

[29] J. Bedford, A. Brandhuber, B. J. Spence, and G. Travaglini, *A Recursion relation for gravity amplitudes*, *Nucl.Phys.* **B721** (2005) 98–110, [`hep-th/0502146`].

[30] A. Hodges, *New expressions for gravitational scattering amplitudes*, `arXiv:1108.2227`.

[31] V. Nair, *A current algebra for some gauge theory amplitudes*, *Phys.Lett.* **B214** (1988) 215.

[32] R. Boels, L. Mason, and D. Skinner, *Supersymmetric Gauge Theories in Twistor Space*, *JHEP* **0702** (2007) 014, [`hep-th/0604040`].

[33] B. Feng and S. He, *Graphs, determinants and gravity amplitudes*, `arXiv:1207.3220`.

[34] C. Cheung, *Gravity Amplitudes from n-Space*, `arXiv:1207.4458`.